## CIRCULATING FAT SOLUBLE VITAMINS AND IONIZED CALCIUM CONCENTRATIONS IN SADDLE-BILLED STORKS (*EPHIPPIORHYNCHUS SENEGALENSIS*) UNDER HUMAN CARE

Kerri A. Slifka, MS1\* and Janis Raines, DVM1

<sup>1</sup>Department of Health, Dallas Zoo, 650 S. RL Thornton FWY, Dallas, TX 75203, USA.

## **Abstract**

Fecundity of saddle-billed storks in human care is dropping from previous levels, and with the report of developmental abnormalities such as fatal malpositions, late term embryo deaths and beak abnormalities in recent hatchlings, there are concerns that there is a nutritional basis for these incidents. Nutritional status of the breeding pair can have a substantial impact on the viability of the offspring. Nutrients transfer from the hen to the egg via the ovary to the yolk or via the oviduct to the albumen, egg shell and membranes. Vitamin A deficiencies have been noted to cause fatal malpositions in chick embryos (Polk & Sipe, 1940). Vitamin D deficiencies have also been implicated in late term embryo death. Beak abnormalities could be caused by a variety of nutritional factors including insufficient vitamin D (Stevens et al., 1984; Schaftenaar & van Leeuwen, 2015) and calcium (Fidgett & Dierenfeld, 2008; Schaftenaar & van Leeuwen, 2015). All North American saddle-billed stork holding institutions were requested to collect blood samples for a vitamin profile (Vitamins A, E, and beta-carotene), and a vitamin D profile (PTH, ionized calcium, 25-hydroxyvitamin D (25-OH D)). Samples were submitted to the Michigan State University Diagnostic Center for Population and Animal Health for analysis. Table 1 presents mean blood values from the twelve responding institutions. As expected, there is substantial variation both within and among zoos. Mean vitamin A and 25-OH D were not significantly different among institutions or sexes. Vitamin E and ionized calcium values were significantly higher in females compared to males across all institutions (P<0.05). Ionized calcium concentrations differed significantly among facilities (P < 0.01). Links with diet and reproduction will be examined in the future.

## Acknowledgements

The authors would like to thank the following institutions for providing sample results:

Albuquerque BioPark Zoo, Assiniboine Park Zoo, The Maryland Zoo in Baltimore, Columbus Zoo, Dallas Zoo, Detroit Zoo, Fort Worth Zoo, Gladys Porter Zoo, Nashville Zoo, North Carolina Zoo, San Diego Zoo, Zoo Miami.

Special thanks to Chase LaDue, M.S. for assistance with statistics.

## Literature Cited

Fidgett AL and Dierenfeld ES (2008) Minerals and stork nutrition. In: M Fowler and RE Miller, Eds. Zoo and Wild Animal Medicine: Current Therapy Vol 6. pp 206-213.

Polk HD and Sipe GR (1940) The effect of vitamin A deficiency on malposition of the chick embryo. *Poult Sci* 19:396-400.

Schaftenaar W and van Leeuwen JP (2015) The influence of ultraviolet-B radiation on the growth of marabou stork (*Leptoptilos crumeniferus*) nestlings in relation to plasma calcium, phosphorus, and vitamin D<sub>3</sub> concentrations. *J Zoo Wildl Med* 46(4):682-690.

Stevens VI, Blair R, Salmon RE, and Stevens JP (1984) Effect of varying levels of dietary vitamin D<sub>3</sub> on turkey hen egg production, fertility, hatchability, embryo mortality and incidence of embryo malformations. *Poult Sci* 63:760-764.

Table 1. Serum Vitamin A, E, 25-OH D and ionized calcium in saddle-billed

storks (Ephippiorhynchus senegalensis) (mean  $\pm$  SEM).

<i>I</i>	(ng/ml) $726.0 \pm 49.2$	$(\mu g/ml)$ 10.61 ± 3.67	(nmol/L)	(mmol/L)
	$726.0 \pm 49.2$	$10.61 \pm 3.67$		
)		10.01 = 5.07	$11.3 \pm 0.3$	$1.40 \pm 0.02$
			$16.5 \pm 1.5$	$1.29 \pm 0.12$
2	$932.5 \pm 159.5$	$7.45 \pm 0.88$	$32.0 \pm 1.0$	$1.40 \pm 0.02$
7	$547.3 \pm 90.8$	$0.42 \pm 1.56$	$21.9 \pm 4.7$	$1.47 \pm 0.02$
	531.0	2.94	37.0	1.38
2	$715.5 \pm 137.$		$22.5 \pm 1.5$	$1.41 \pm 0.04$
1	$596.8 \pm 32.3$	(B) A (B)	$19.8 \pm 2.4$	$1.33 \pm 0.03$
2	$667.0 \pm 90.0$	$6.89 \pm 1.01$	$15.0 \pm 3.0$	$1.46 \pm 0.01$
1	596.0	9.71	21.0	1.27
2	$470.5 \pm 32.5$	$10.56 \pm 1.52$	$13.5 \pm 2.5$	$1.33 \pm 0.06$
2	$578.5 \pm 46.5$	$5.24 \pm 0.39$	$45.0 \pm 2.0$	$1.22 \pm 0.04$
_	594.0	11.52	54.0	1.32
1 2 2		$715.5 \pm 137.$ $596.8 \pm 32.3$ $667.0 \pm 90.0$ 596.0 $470.5 \pm 32.5$ $578.5 \pm 46.5$	$715.5 \pm 137.$ $7.20 \pm 0.52$ $596.8 \pm 32.3$ $3.84 \pm 0.32$ $667.0 \pm 90.0$ $6.89 \pm 1.01$ 596.0 $9.71470.5 \pm 32.5 10.56 \pm 1.52578.5 \pm 46.5 5.24 \pm 0.39$	$715.5 \pm 137. \qquad 7.20 \pm 0.52 \qquad 22.5 \pm 1.5$ $596.8 \pm 32.3 \qquad 3.84 \pm 0.32 \qquad 19.8 \pm 2.4$ $667.0 \pm 90.0 \qquad 6.89 \pm 1.01 \qquad 15.0 \pm 3.0$ $596.0 \qquad 9.71 \qquad 21.0$ $470.5 \pm 32.5 \qquad 10.56 \pm 1.52 \qquad 13.5 \pm 2.5$ $578.5 \pm 46.5 \qquad 5.24 \pm 0.39 \qquad 45.0 \pm 2.0$

<sup>&</sup>lt;sup>1</sup>25-hydroxyvitamin D